Supercongruences between truncated ${}_3F_2$ hypergeometric series

Ji-Cai Liu

Department of Mathematics, East China Normal University, Shanghai 200241, PR China
jc2051@163.com

arXiv:1702.07232v1 [math.NT] 19 Feb 2017

Abstract. We establish four supercongruences between truncated ${}_3F_2$ hypergeometric series involving p-adic Gamma functions, which extend some of the Rodriguez-Villegas supercongruences.

Keywords: Supercongruences; Truncated hypergeometric series; p-adic Gamma functions

MR Subject Classifications: Primary 11A07; Secondary 33C20

1 Introduction

For complex numbers a_i, b_j and z, with none of the b_j being negative integers or zero, the truncated hypergeometric series are defined as

$$ {}_rF_s\left[\begin{matrix}a_1, a_2, \cdots, a_r\\ b_1, b_2, \cdots, b_s\end{matrix}; z\right]_n = \sum_{k=0}^{n-1} \frac{(a_1)_k (a_2)_k \cdots (a_r)_k}{(b_1)_k (b_2)_k \cdots (b_s)_k} \cdot \frac{z^k}{k!},$$

where $(a)_0 = 1$ and $(a)_k = a(a+1)\cdots(a+k-1)$ for $k \geq 1$.

Throughout this paper, let $p \geq 5$ be a prime. Rodriguez-Villegas [13] conjectured 22 supercongruences to establish the relationship between the number of points over F_p on certain Calabi-Yau manifolds and truncated hypergeometric series. In particular, he conjectured four supercongruences on truncated ${}_3F_2$ hypergeometric series, which relate to certain modular K3 surfaces. These were all of the form

$$ {}_3F_2\left[\begin{matrix}\frac{1}{2}, -a, a+1\\ 1, 1\end{matrix}; 1\right]_p \equiv c_p \pmod{p^2},$$

where $a = -\frac{1}{2}, -\frac{1}{3}, -\frac{1}{4}, -\frac{1}{6}$ and c_p is the p-th Fourier coefficient of a weight three modular form on a congruence subgroup of $SL(2,\mathbb{Z})$.

The case when $a = -\frac{1}{2}$ was originally conjectured by Beukers and Stienstra [2], and confirmed by van Hamme [17], Ishikawa [5] and Ahlgren [1]. The other cases when $a = -\frac{1}{3}, -\frac{1}{4}, -\frac{1}{6}$ were partially proved by Mortenson [12], and finally proved by Z.-W. Sun [15].

Recall that the p-adic Gamma function [3, Definition 11.6.5] is defined as

$$\Gamma_p(x) = \lim_{m \to x} (-1)^m \prod_{\substack{1<k<m\\(k,p)=1}} k,$$

where the limit is for m tending to x p-adically in $\mathbb{Z}_{\geq 0}$. Let $\langle a\rangle_p$ denote the least non-negative integer r with $a \equiv r \pmod{p}$.

Z.-H. Sun [14, Theorem 2.5] and the author [10, Theorem 1.2] proved that for any p-adic integer a,

$$\begin{aligned}&{}_3F_2\left[\begin{matrix}\frac{1}{2}, -a, a+1\\ 1, 1\end{matrix}; 1\right]_p\\ &\quad\equiv \begin{cases}0, & \text{if } \langle a\rangle_p \equiv 1 \pmod 2\\ (-1)^{\frac{p+1}{2}}\Gamma_p\left(-\frac{a}{2}\right)^2\Gamma_p\left(\frac{a+1}{2}\right)^2, & \text{if } \langle a\rangle_p \equiv 0 \pmod 2\end{cases} \pmod{p^2}, \qquad (1.1)\end{aligned}$$

which extends the above four Rodriguez-Villegas supercongruences. Guo and Zeng [4, Theorem 1.3] have established an interesting q-analogue of the first case of (1.1).

Recently, the author [9] has extended the first case of (1.1) and showed that for any positive integer n and p-adic integer a with $\langle a\rangle_p \equiv 1 \pmod 2$,

$${}_3F_2\left[\begin{matrix}\frac{1}{2}, -a, a+1\\ 1, 1\end{matrix}; 1\right]_{np} \equiv 0 \pmod{p^2}.$$

In this paper, we aim to prove the following supercongruences between truncated ${}_3F_2$ hypergeometric series.

Theorem 1.1 *Let $p \geq 5$ be a prime and n be a positive integer. For $a \in \left\{-\frac{1}{2}, -\frac{1}{3}, -\frac{1}{4}, -\frac{1}{6}\right\}$ with $\langle a\rangle_p \equiv 0 \pmod 2$, we have*

$$\begin{aligned}&{}_3F_2\left[\begin{matrix}\frac{1}{2}, -a, a+1\\ 1, 1\end{matrix}; 1\right]_{np}\\ &\quad\equiv (-1)^{\frac{p+1}{2}}\Gamma_p\left(-\frac{a}{2}\right)^2\Gamma_p\left(\frac{a+1}{2}\right)^2 {}_3F_2\left[\begin{matrix}\frac{1}{2}, -a, a+1\\ 1, 1\end{matrix}; 1\right]_n \pmod{p^2}. \qquad (1.2)\end{aligned}$$

Replacing n by p^{r-1} in (1.2) and then using induction, we immediately get the following results:

Corollary 1.2 *Let $p \geq 5$ be a prime and r be a positive integer. For $a \in \left\{-\frac{1}{2}, -\frac{1}{3}, -\frac{1}{4}, -\frac{1}{6}\right\}$ with $\langle a\rangle_p \equiv 0 \pmod 2$, we have*

$${}_3F_2\left[\begin{matrix}\frac{1}{2}, -a, a+1\\ 1, 1\end{matrix}; 1\right]_{p^r} \equiv (-1)^{\frac{r(p+1)}{2}}\Gamma_p\left(-\frac{a}{2}\right)^{2r}\Gamma_p\left(\frac{a+1}{2}\right)^{2r} \pmod{p^2}.$$

Numerical calculation suggests that supercongruence (1.2) cannot be extended to any p-adic integer a with $\langle a\rangle_p \equiv 0 \pmod 2$ in the direction of (1.1). In the next section, we first recall some lemmas that we will need later. The proof of Theorem 1.1 will be given in the last section.

2 Some lemmas

The Fermat quotient of an integer a with respect to an odd prime p is given by

$$q_p(a) = \frac{a^{p-1}-1}{p}.$$

Lemma 2.1 *(Eisenstein) Suppose p is an odd prime and r is a positive integer. For non-zero p-adic integers a and b, we have*

$$q_p(ab) \equiv q_p(a) + q_p(b) \pmod{p},$$
$$q_p(a^r) \equiv r q_p(a) \pmod{p}.$$

Lemma 2.2 *(Lehmer [6]) Let $H_n = \sum_{k=1}^{n} \frac{1}{k}$ be the n-th harmonic number. For any prime $p \geq 5$, we have*

$$H_{\lfloor p/2 \rfloor} \equiv -2q_p(2) \pmod{p},$$

where $\lfloor x \rfloor$ denotes the greatest integer less than or equal to a real number x.

Lemma 2.3 *([8, Lemma 2.4]) Let $p \geq 5$ be a prime, r and $0 \leq k \leq p-1$ be non-negative integers. For $a \in \left\{-\frac{1}{2}, -\frac{1}{3}, -\frac{1}{4}, -\frac{1}{6}\right\}$, we have*

$$\frac{(-a)_{k+rp}(a+1)_{k+rp}}{(1)_{k+rp}^2} \equiv \frac{(-a)_r(a+1)_r}{(1)_r^2} \cdot \frac{(-a)_k(a+1)_k}{(1)_k^2} \times \left(1 + 2rpH_{\lfloor -pa \rfloor} - 2rpH_k + rp\sum_{i=0}^{k-1}\left(\frac{1}{-a+i} + \frac{1}{a+1+i}\right)\right) \pmod{p^2}. \tag{2.1}$$

Lemma 2.4 *([7, Lemma 2.1]) Suppose $p \geq 5$ is a prime, r and $0 \leq k \leq p-1$ are non-negative integers. Then*

$$\binom{2rp+2k}{rp+k} \equiv \binom{2r}{r}\binom{2k}{k}(1 + 2rp(H_{2k} - H_k)) \pmod{p^2}. \tag{2.2}$$

Lemma 2.5 *([10, (2.9) & (2.18)] and [16, (16) & (17)]) For any even integer n, we have*

$$\sum_{k=0}^{n}\binom{2k}{k}^2\binom{n+k}{2k}\left(-\frac{1}{4}\right)^k = \frac{\binom{n}{n/2}^2}{4^n}, \tag{2.3}$$

$$\sum_{k=0}^{n}\binom{2k}{k}^2\binom{n+k}{2k}\left(-\frac{1}{4}\right)^k H_k = \frac{\binom{n}{n/2}^2}{4^n}H_n, \tag{2.4}$$

$$\sum_{k=0}^{n}\binom{2k}{k}^2\binom{n+k}{2k}\left(-\frac{1}{4}\right)^k H_{2k} = \frac{\binom{n}{n/2}^2}{2\cdot 4^n}H_n, \tag{2.5}$$

$$\sum_{k=0}^{n}\binom{2k}{k}^2\binom{n+k}{2k}\left(-\frac{1}{4}\right)^k \sum_{i=0}^{k-1}\frac{1}{n+1+i} = \frac{\binom{n}{n/2}^2}{4^n}\left(\frac{3}{2}H_n - H_{n/2}\right). \tag{2.6}$$

Lemma 2.6 *Let $p \geq 5$ be a prime. For $a \in \left\{-\frac{1}{2}, -\frac{1}{3}, -\frac{1}{4}, -\frac{1}{6}\right\}$, we have*

$$H_{\lfloor -pa \rfloor} \equiv H_{\langle a \rangle_p} \pmod{p}. \tag{2.7}$$

Proof. For any prime $p \geq 5$, there exists $\varepsilon \in \{1, -1\}$ such that $p \equiv \varepsilon \pmod{2,3,4,6}$. We give the proof of (2.7) for $a = -\frac{1}{3}$. The proofs of the other three cases run similarly.

If $p \equiv 1 \pmod 3$, then $\lfloor p/3 \rfloor = (p-1)/3$ and $\langle -1/3 \rangle_p = (p-1)/3$, and so (2.7) clearly holds for $a = -\frac{1}{3}$.

If $p \equiv -1 \pmod 3$, then $\lfloor p/3 \rfloor = (p-2)/3$ and $\langle -1/3 \rangle_p = (2p-1)/3$. Applying the fact that $H_k = H_{p-1-k} \pmod p$ for $0 \leq k \leq p-1$, we conclude that (2.7) also holds for $a = -\frac{1}{3}$. □

3 Proof of Theorem 1.1

We first prove that for any non-negative integer r,

$$\begin{aligned}
&\sum_{k=rp}^{(r+1)p-1} \frac{\left(\frac{1}{2}\right)_k (-a)_k (a+1)_k}{(1)_k^3} \\
&\qquad \equiv (-1)^{\frac{p+1}{2}} \Gamma_p\left(-\frac{a}{2}\right)^2 \Gamma_p\left(\frac{a+1}{2}\right)^2 \cdot \frac{\left(\frac{1}{2}\right)_r (-a)_r (a+1)_r}{(1)_r^3} \pmod{p^2}.
\end{aligned} \tag{3.1}$$

Noting that

$$\frac{\left(\frac{1}{2}\right)_k (-a)_k (a+1)_k}{(1)_k^3} = \binom{2k}{k}^2 \binom{a+k}{2k} \left(-\frac{1}{4}\right)^k, \tag{3.2}$$

and letting $k \to k + rp$ on the left-hand side of (3.1), we see that (3.1) is equivalent to

$$\begin{aligned}
&\sum_{k=0}^{p-1} \binom{2k+2rp}{k+rp}^2 \binom{a+k+rp}{2k+2rp} \left(-\frac{1}{4}\right)^{k+rp} \\
&\qquad \equiv (-1)^{\frac{p+1}{2}} \Gamma_p\left(-\frac{a}{2}\right)^2 \Gamma_p\left(\frac{a+1}{2}\right)^2 \binom{2r}{r}^2 \binom{a+r}{2r} \left(-\frac{1}{4}\right)^r \pmod{p^2}.
\end{aligned} \tag{3.3}$$

By Lemma 2.1 and 2.2, we have

$$\begin{aligned}
\left(\frac{1}{4}\right)^{k+rp} &= \left(\frac{1}{4}\right)^{k+r} \cdot \left(\frac{1}{4}\right)^{r(p-1)} \\
&= \left(\frac{1}{4}\right)^{k+r} \left(1 + pq_p\left(4^{-r}\right)\right) \\
&\equiv \left(\frac{1}{4}\right)^{k+r} \left(1 - 2rpq_p(2)\right) \\
&\equiv \left(\frac{1}{4}\right)^{k+r} \left(1 + rpH_{\lfloor p/2 \rfloor}\right) \pmod{p^2}.
\end{aligned} \tag{3.4}$$

Combining (2.2) and (3.4) gives

$$\left(\frac{1}{4}\right)^{k+rp}\binom{2k+2rp}{k+rp}$$
$$\equiv\left(\frac{1}{4}\right)^{k+r}\binom{2r}{r}\binom{2k}{k}(1+rp(2H_{2k}-2H_k+H_{\lfloor p/2\rfloor}))\pmod{p^2}. \tag{3.5}$$

On the other hand, using (2.7) and

$$\frac{(-a)_k(a+1)_k}{(1)_k^2}=(-1)^k\binom{2k}{k}\binom{a+k}{2k},$$

we can rewrite (2.1) as

$$(-1)^{k+rp}\binom{2k+2rp}{k+rp}\binom{a+k+rp}{2k+2rp}$$
$$\equiv(-1)^{r+k}\binom{2r}{r}\binom{a+r}{2r}\binom{2k}{k}\binom{a+k}{2k}$$
$$\times\left(1+2rpH_{\langle a\rangle_p}-2rpH_k+rp\sum_{i=0}^{k-1}\left(\frac{1}{-a+i}+\frac{1}{a+1+i}\right)\right)\pmod{p^2}. \tag{3.6}$$

Applying (3.5) and (3.6) to the left-hand side of (3.3) yields

$$\text{LHS (3.3)}$$
$$\equiv\binom{2r}{r}^2\binom{a+r}{2r}\left(-\frac{1}{4}\right)^r\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{a+k}{2k}\left(-\frac{1}{4}\right)^k$$
$$\times\left(1+rp\left(2H_{2k}-4H_k+2H_{\langle a\rangle_p}+H_{\lfloor p/2\rfloor}+\sum_{i=0}^{k-1}\left(\frac{1}{-a+i}+\frac{1}{a+1+i}\right)\right)\right)\pmod{p^2}. \tag{3.7}$$

In fact, the proof of (3.3) follows from the following congruence:

$$\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{a+k}{2k}\left(-\frac{1}{4}\right)^k$$
$$\times\left(2H_{2k}-4H_k+2H_{\langle a\rangle_p}+H_{\lfloor p/2\rfloor}+\sum_{i=0}^{k-1}\left(\frac{1}{-a+i}+\frac{1}{a+1+i}\right)\right)\equiv 0\pmod{p}. \tag{3.8}$$

Substituting (3.8) into (3.7), and then using (3.2) and (1.1), we obtain

$$\text{LHS (3.3)}\equiv\binom{2r}{r}^2\binom{a+r}{2r}\left(-\frac{1}{4}\right)^r\cdot{}_3F_2\left[\begin{matrix}\frac{1}{2},-a,a+1\\1,1\end{matrix};1\right]_p$$
$$\equiv(-1)^{\frac{p+1}{2}}\Gamma_p\left(-\frac{a}{2}\right)^2\Gamma_p\left(\frac{a+1}{2}\right)^2\binom{2r}{r}^2\binom{a+r}{2r}\left(-\frac{1}{4}\right)^r\pmod{p^2},$$

which is (3.3). So it suffices to prove (3.8).

Since $a \equiv \langle a\rangle_p \pmod p$, we have

$$\begin{aligned}&\text{LHS (3.8)}\\ &\equiv \sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{\langle a\rangle_p+k}{2k}\left(-\frac{1}{4}\right)^k\\ &\times\left(2H_{2k}-4H_k+2H_{\langle a\rangle_p}+H_{\lfloor p/2\rfloor}+\sum_{i=0}^{k-1}\left(\frac{1}{-\langle a\rangle_p+i}+\frac{1}{\langle a\rangle_p+1+i}\right)\right)\pmod p.\end{aligned}\tag{3.9}$$

By (2.3)-(2.6) and the fact that $0\le\langle a\rangle_p\le p-1$ is even, we obtain

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{\langle a\rangle_p+k}{2k}\left(-\frac{1}{4}\right)^k\left(2H_{2k}-4H_k+2H_{\langle a\rangle_p}+H_{\lfloor p/2\rfloor}+\sum_{i=0}^{k-1}\frac{1}{\langle a\rangle_p+1+i}\right)\\ &=\frac{\binom{\langle a\rangle_p}{\langle a\rangle_p/2}^2}{4^{\langle a\rangle_p}}\left(H_{\lfloor p/2\rfloor}+\frac{1}{2}H_{\langle a\rangle_p}-H_{\langle a\rangle_p/2}\right).\end{aligned}\tag{3.10}$$

Let $b=p-\langle a\rangle_p$. It is clear that $\langle a\rangle_p\equiv -b \pmod p$ and $0\le b-1\le p-1$ is even. Using (2.6) and the fact that $\binom{-b+k}{2k}=\binom{b-1+k}{2k}$, we get

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{\langle a\rangle_p+k}{2k}\left(-\frac{1}{4}\right)^k\sum_{i=0}^{k-1}\frac{1}{-\langle a\rangle_p+i}\\ &\equiv\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{-b+k}{2k}\left(-\frac{1}{4}\right)^k\sum_{i=0}^{k-1}\frac{1}{b+i}\pmod p\\ &=\sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{b-1+k}{2k}\left(-\frac{1}{4}\right)^k\sum_{i=0}^{k-1}\frac{1}{b+i}\\ &=\frac{\binom{b-1}{(b-1)/2}^2}{4^{b-1}}\left(\frac{3}{2}H_{b-1}-H_{(b-1)/2}\right).\end{aligned}\tag{3.11}$$

By (2.3), we have

$$\begin{aligned}\frac{\binom{b-1}{(b-1)/2}^2}{4^{b-1}} &= \sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{b-1+k}{2k}\left(-\frac{1}{4}\right)^k\\ &\equiv \sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{-\langle a\rangle_p-1+k}{2k}\left(-\frac{1}{4}\right)^k \pmod p\\ &= \sum_{k=0}^{p-1}\binom{2k}{k}^2\binom{\langle a\rangle_p+k}{2k}\left(-\frac{1}{4}\right)^k\\ &= \frac{\binom{\langle a\rangle_p}{\langle a\rangle_p/2}^2}{4^{\langle a\rangle_p}}. \end{aligned}\tag{3.12}$$

Substituting (3.10)-(3.12) into (3.9) gives

$$\begin{aligned}\text{LHS (3.8)} \equiv{}& \frac{\binom{\langle a\rangle_p}{\langle a\rangle_p/2}^2}{4^{\langle a\rangle_p}}\\ &\times\left(H_{\lfloor p/2\rfloor}+\frac{1}{2}H_{\langle a\rangle_p}-H_{\langle a\rangle_p/2}+\frac{3}{2}H_{p-1-\langle a\rangle_p}-H_{(p-1-\langle a\rangle_p)/2}\right) \pmod p.\end{aligned}\tag{3.13}$$

Note that

$$\begin{aligned}H_{(p-1-\langle a\rangle_p)/2} &= H_{(p-1)/2}-\sum_{i=0}^{\langle a\rangle_p/2-1}\frac{1}{(p-1)/2-i}\\ &\equiv H_{\lfloor p/2\rfloor}-\sum_{i=0}^{\langle a\rangle_p/2-1}\frac{1}{-1/2-i} \pmod p\\ &= H_{\lfloor p/2\rfloor}+2H_{\langle a\rangle_p}-H_{\langle a\rangle_p/2}.\end{aligned}$$

Substituting the above congruence into (3.13) yields

$$\text{LHS (3.8)} \equiv \frac{3}{2}\cdot\frac{\binom{\langle a\rangle_p}{\langle a\rangle_p/2}^2}{4^{\langle a\rangle_p}}\left(H_{p-1-\langle a\rangle_p}-H_{\langle a\rangle_p}\right)\equiv 0 \pmod p,$$

where we have utilized the fact that $H_{p-1-k}\equiv H_k \pmod p$ for $0\le k\le p-1$. This concludes the proof of (3.8).

Taking the sum over r from 0 to $n-1$ on both sides of (3.1) gives

$$\begin{aligned}&{}_3F_2\left[\begin{matrix}\frac{1}{2},-a,a+1\\1,1\end{matrix};1\right]_{np}\\ &\quad\equiv(-1)^{\frac{p+1}{2}}\Gamma_p\left(-\frac{a}{2}\right)^2\Gamma_p\left(\frac{a+1}{2}\right)^2{}_3F_2\left[\begin{matrix}\frac{1}{2},-a,a+1\\1,1\end{matrix};1\right]_n \pmod{p^2}.\end{aligned}$$

This completes the proof of Theorem 1.1.

References

[1] S. Ahlgren, Gaussian hypergeometric series and combinatorial congruences, Symbolic computation, number theory, special functions, physics and combinatorics (Gainesville, FL, 1999), 1–12, Dev. Math., 4, Kluwer, Dordrecht, 2001.

[2] F. Beukers and J. Stienstra, On the Picard-Fuchs equation and the formal Brauer group of certain elliptic K3 surfaces, Math. Ann. 271 (1985), 269–304.

[3] H. Cohen, Number Theory. Vol. II. Analytic and Modern Tools, Grad. Texts in Math., vol. 240, Springer, New York, 2007.

[4] V.J.W. Guo and J. Zeng, Some q-supercongruences for truncated basic hypergeometric series, Acta Arith. 171 (2015), 309–326.

[5] T. Ishikawa, Super congruence for the Apéry numbers, Nagoya Math. J. 118 (1990), 195–202.

[6] E. Lehmer, On congruences involving Bernoulli numbers and the quotients of Fermat and Wilson, Ann. Math. 39 (1938), 350–360.

[7] J.-C. Liu, On two conjectural supercongruences of Apagodu and Zeilberger, J. Difference Equ. Appl. 22 (2016), 1791–1799.

[8] J.-C. Liu, Congruences for truncated hypergeometric series ${}_2F_1$, Bull. Aust. Math. Soc., in press. (arXiv:1605.06632)

[9] J.-C. Liu, Some supercongruences for truncated generalized hypergeometric series ${}_3F_2$, preprint, 2016, arXiv:1610.03799.

[10] J.-C. Liu, Generalized Rodriguez-Villegas supercongruences involving p-adic Gamma functions, preprint, 2016, arXiv:1611.07686.

[11] L. Long and R. Ramakrishna, Some supercongruences occurring in truncated hypergeometric series, Adv. Math. 290 (2016), 773–808.

[12] E. Mortenson, Supercongruences for truncated ${}_{n+1}F_n$ hypergeometric series with applications to certain weight three newforms, Proc. Amer. Math. Soc. 133 (2005), 321–330.

[13] F. Rodriguez-Villegas, Hypergeometric families of Calabi-Yau manifolds, in: Calabi-Yau Varieties and Mirror Symmetry (Toronto, ON, 2001), Fields Inst. Commun., 38, Amer. Math. Soc., Providence, RI, 2003, 223–231.

[14] Z.-H. Sun, Generalized Legendre polynomials and related supercongruences, J. Number Theory 143 (2014), 293–319.

[15] Z.-W. Sun, On sums involving products of three binomial coefficients, Acta Arith. 156 (2012), 123–141.

[16] R. Tauraso, Supercongruences related to ${}_3F_2(1)$ involving harmonic numbers, preprint, 2017, arXiv:1701.00729.

[17] L. van Hamme, Proof of a conjecture of Beukers on Apéry numbers, Proceedings of the conference on p-adic analysis (Houthalen, 1987), 189–195.